

PUBLIC HEALTH IMPLICATIONS OF LOW BIRTHWEIGHT*

Leonardo Mata", JJJn J. Urrutia"· and Edgar Mohs***.

Institute of Investigations in Health (INISA), University of Costa Rica
 and Ministry of Health of Costa Rica

Institute of Nutrition of Central America and Panama (INCAPI,
 Guatemala, C. A.

National Children'S Hospital, Caja Costarricense del Seguro Social (CCSS),
 Costa Rica

SUMMARY

The present paper interprets the significance of anthropometric data of newborns from a typical Indian village of Guatemala and from the general population of Costa Rica. A good correlation exists between a deficient fetal growth and infant mortality and retarded physical growth. The significance of fetal growth retardation on early and late malnutrition and defects is discussed.

The existing information reveals a high incidence of neonates with altered fetal growth and development in Latin America. This circumstance is the main factor associated with the high infant mortality and retarded development observed in many populations of the Subcontinent.

~ Presented at: Colloquium on "Prenatal and Perinatal Nutrition", held as part of the IV Latin American Nutrition Congress, Caracas, Venezuela, 21-27 November 1976, under the sponsorship of the Latin American Nutrition Society (SLAN).

*** Director of the above-mentioned Institute and Adviser to the Ministry of Health of Costa Rica.

**** Member of the Division of Biology and Human Nutrition of INCAP.

..... Member of the Hospital, CCSS, San Jose, Costa Rica.

Certain countries have attained a substantial improvement in the conditions affecting fetal growth, by implementing comprehensive health and development programs with a holistic approach. The interventions that need to be strengthened in order to improve fetal development in our nations are reviewed.

I. INTRODUCTION

Knowledge of the incidence and causality of low birth weight is of singular importance for the formulation of health policies and programs, especially when interventions of great coverage and extension are planned. This condition is particularly true for the developing countries, since in them fetal growth retardation and premature delivery occur with greater frequency than in the industrialized countries.

As was to be expected, practically no information is available on fetal growth in the developing countries, especially for suburban and rural populations. In general, the existing information is deficient in quality and not representative of the total population. Even in the case of urban areas, information is also scarce and not representative.

These considerations explain why sufficient importance is not given to low birth weight as a health problem in developing countries.

II. STUDY MATERIAL

To illustrate the implications of low birth weight on health, data derived from a prospective study (The Cauque Study) of a rural community of Guatemala will be employed,¹⁻³ as well as data from the Instituto Materno-Infantil CARIT (Mother and Child Health Institute CARIT) and from the General Direction of Statistics and Census, both in Costa Rica.⁴⁻⁵ The rural community of Guatemala is located at an altitude of 2,000 m above sea level and belongs to the linguistic Maya-Cakchiquel group. During the 1964-1973 period careful information was collected for practically the totality of newborn cohorts; these were observed in a prospective form. As a whole, the population consisted of 430 liveborn infants who represented more than 95% of the total births. Among many other determinations,³ studies included measuring in a prospective way the weight and height, and survival of all the children.

The Costa Rica data correspond to a random sample of 10% of all liveborns in certain years at the CARIT Institute. The study period comprised from 1965 to 1974. During these years, this hospital center provided medical attention to a population representative of all social sectors of the country, including the rural population. Information on infant mortality and birth rate in Costa Rica corresponds to the total population of the country in the stated period.

III. LOW BIRTH WEIGHT IN THE DEVELOPING COUNTRIES

Available information in Latin America reveals a serious fetal growth problem.^{6,8} For example, the mean birth weight in the Guatemalan village (Table 0,9 is one of the lowest registered in the literature and is significantly under that observed in industrialized populations located at similar altitudes than those: of the Guatemalan highlands,¹⁰ as well as of Indian groups with better economic situation.¹¹ Furthermore, the fact that the anthropometric characteristics of the newborn in the Indian village studied did not appear to change during the 10-year period of observation, ..., is suggestive that the factors which determine a deficient fetal growth remained constant during such period.

TABLE 1

WEIGHT OF LIVE NEWBORN COHORTS,
SANTA MARIA CAUaUE, GUATEMALA

Ye.r	No. of children	Weight,g Average ± D.E.	0/0 < 2,501 g
1984	37	2,595 ± 360	35
1966	46	2,506 ± 321	46
1968	67	2,510 ± 422	44
1970	67	2,558 ± 412	36
1971	60	2,564 ± 328	48

Classification of the 415 neonates by weight at birth and by gestational age showed that 7% were pre-term and that an additional 340/0 were term-small-for-gestational age. The ratio "pre-term: term-small-for-gestational age" which was of 1:4.7 in the village contrasts with that observed in more developed societies, where it is generally of 2:1 (Table 2).

A high incidence of low birth weight has been described in pre-industrialized populations, including Latin America].^{8, 12-18} Nevertheless, truly representative data for this Region do not exist, particularly in what concerns rural areas; in addition there is no information on fetal maturity due to the difficulties inherent to the collection of information on gestational age.

TABLE 2
FETAL MATURITY OF 415 NEONATES,
SANTA MARIA CAUQUE, 1964-1972

Classification	No_ a/a	Birth weight, g	Gestational age, weeki
Term	43 (10.3)	> 3,000	37 -42 ~
	199 (47.8)	2,501-3,000	37-42
	143 (34.4)	< 2,501	37-42
Pre-term	6 (1.4)	2,001-2.500	35 -37
	24 (5.8'	< 2,001	31-36

IV. CORRELATION OF LOW BIRTH WEIGHT WITH MORTALITY

The influence of low birth weight and fetal immaturity on the neonatal mortality has been amply demonstrated.¹⁹⁻²³ In the Cauque Study the correlation between birth weight and infant mortality (Table 3) was so marked as to characterize said variable as the best predictive survival index in similar naral populations: Actually, the indicator is closely correlated with gestational age (Table 4) which also showed a strong association with survival. A birth weight of at least 2,750 g and a gestational age of more than 36 weeks was associated with absolute survival during the neonatal period, and with a relatively good survival during the rest of the flJ'st year of life. It must be taken into account that the totality of the neonates who survived the first 48 hours were adequately breast-fed for long periods of time. Definitive weaning takes place during the second or third years of life.³ Introduction of liquid foods starts at the age of 3 to 6 months, and of solid foods, at 6 to 12 months.

The combination of both variables has been proposed by several authors²⁴⁻²⁶ and is shown in Table 5 for the Cauque Study. The pre-term infants constituted the group with the highest and sustained mortality during all the course of their first year of life. However, the surviving babies showed a high survival during the rest of their preschool period.

The born at-term infants but small for their gestational age (malnourished fetuses) manifested a high mortality during all of the flJ'st and second years of life, as well as a high mortality during the rest of the preschool age.

Infants born at term but weighing more than 2,500 g exhibited the lowest neonatal mortality (8 per 1,000) and a post-neonatal infant mortality also relatively low. However, many of these were vulnerable during the preschool years, resulting in a high age-specific mortality in each one of the three preschool years.

Ct

TABLE 3

INFANT MORTALITY AND BIRTH WEIGHT, SANTA MARIA CAUQUE,
GUATEMALA, 1964-1173

Birth weight. g	No. of children	Age			< 1 year
		<29 days	29 days-5 months	6-11 months	
< 1.501	5	31600)	1 (200)	0	41800)
1,601-1,750	11	2 (182)	3 (273)	1 (91)	6 (545)
1,751-2.000	17	4 (235)	4 (235)	1 (59)	9 (529)
2,001-2.250	47	2 (43)	2 (43)	2 (43)	6 (128)
2,251-2,500	99	3 (301)	0	1 (10)	4 (40)
2,501-2.750	125	2 16)	3 24)	21161	7(56)
2,751-3,000	82	0	2 24)	2 (24)	4 (49)
t) 3,001-3,250	32	0	0	0	0
3.251-3,500	11	0	0	1 (91)	1 (91)
(3.501	1	0	0	0	0
rota	430	16 (37)	15 351	10 (23)	41 95)

^{tt} figures in parentheses represent deaths and mortality per 1,000 liveborns of said weight.

TABLE 4

INFANT MORTALITY AND GESTATIONAL AGE, SANTA MARIA CAUQUE,
GUATEMALA, 19M-1973

Gestational ag., weeks	No. of children	Age			
		< 29 days	29 days- 5 months	6-11 months	< 1 year ~ ~
31-32	3	2 (667)	0	0	2 (667)
33-34	8	3 (375)	3 (375)	0	6 (750)
35-36	20	5 (250)	1 (50)	2 (100)	8 (400)
37-38	47	0	2 (43)	a	2 (43)
39-40	261	6 (23)	7 (27)	8 (31)	21 (80)
41-42	77	0	1 (13)	0	1 (13)
Total	416	16 (38)	14 (34)	10 (24)	40 (96)

~ Figures in parentheses represent deaths and mortality per 1,000 livebirths of
Mid gestatio.,.1 age_

TABLES

MORTALITY BY FETAL MATURITY, 416 COHORT CHILDREN,
SANTA MARIA CAUQUE, GUATEMALA, 1964-'972

Classification	1st year			2nd year	3rd year	4th year
	<29 days	29 days. 11 months	<'lt year			
Pre-term > 37 9	101323' 31	6(286) 21	16(516) 31	0 15	0 15	0 8
Term < 2,501 g	4128' 143	8(58) 139	12(84) 143	8(76) 105	3(39, 78	3(50) 60
Term > 2.500 9	2(81) 242	10(42) 240	12(50) 242	9(441 204	5(33) 153	1(8) 122
Total	16(39) 416	24(601 400	40(96} 416	17(52) 324	8(33) 244	4(21) 190

* Figures in parentheses represent deaths and mortality per 1,000 livebirths at the initiation of the period. The underlined figures under mortality rate indicate number of children in cohort at initiation of period.

The importance of an adequate fetal development to avoid neonatal deaths is obvious when this is compared with infant mortality in two contrasting populations: one of them, the Indian village, and the other a population of the East Coast of the United States of America (Table 6).²⁷ No differences are noted in neonatal mortality between the Indian community and the North American population if one takes into account the weight category at birth. Consequently, the high neonatal mortality rates observed in the Indian community and in the developing societies are in great measure due to the higher incidence of low birth weight. Rates higher than those observed in the Cauque Study may be due to the joint action of a deficient diet (particularly during early weaning) and infections.^{3~ 27-29}

TABLE 6
COMPARISON OF INFANT MORTALITY OF TWO POPULATIONS,
BY WEIGHT AT BIRTH

Birth weight, g	Neonatal			Postneonatal			Infantile		
	S.M.C.*	B.	R.**	S.M.C.	B.	R.	S.M.C.	B.	R.
1501-2000	273	210	1.3	303	26	11.7	576	199	2.9
2001-2500	34	45	0.8	34	13	2.6	68	54	1.3
2501-3000	10	10	1.0	43	7	6.1	53	17	3.1
3001-3500	0	5		23	5	4.6	23	10	2.3

* S.M.C. = Santa Marr. Cauq", Guatemala; B. = BaltimOfe, USA.46

** S.M.C./B ratio.

V. RELATIONSHIP OF LOW BIRTH WEIGHT WITH PHYSICAL GROWTH

The Cauque Study demonstrated a marked weight and height deficit of the Indian population with respect to accepted standards.¹⁻³ Other authors have described similar deficiencies.^{30, 31} The Cauque Study also revealed that physical growth deficiencies correlated with the fetal growth.¹⁻³ Birth weight and gestational age, or both variables combined, showed a good correlation with physical growth of the lactating and preschool child (body weight, height and head circumference). Children with low birth weight exhibited a dear tendency to remain in the lower weight tracks (Figure 1) while those with better birth weight occupied the upper growth tracks.

Correlation with gestational age also was high, but only two populations were noted separated by the limit of 36-37 gestational weeks (Figure 2).

As was to be expected, with regards to fetal maturity, the at-term infants with a birth weight of more than 3,000 g grew better than any other group; the pre-term infants weighing less than 2,001 g grew the worst (Figure 3). However, the pre-term group of children with a birth weight greater than 2,000 g grew as well as the term children.¹

The relation between birth weight and physical growth has been explored by other authors through retrospective and prospective analyses, who found a positive correlation between both.³²⁻³³ The few prospective investigations carried out in industrialized countries showed an abnormal growth on the part of pre-term and




Fig. 1. Weight curves (± 2 E. E.) of foUT cohorts of children defined by birth weight in comparison with the mean of the foUT standard curve. Prospective study, Santa Maria Cauque, Guatemala. 1964-1972. Figures in curves are number of children measured²




Fig. 3 *Boertje weight curves of four cohorts of children defined by fetal maturity at birth, in comparison with the mean of the Iowa standard curve. Prospective study, Santa Maria Cauque, Guatemala, 1964-1972.1*

small-for-gestational age infants,^{34, 35} despite the fact that the environment where these children developed will more favorable than is to be expected in developing countries, particularly in rural areas.

There are no similar data to those of the Cauque Study for other developing populations, but a follow-up of Nigerian and Gambian infants showed a deficient weight gain in those with low birth weight.³⁶⁻³⁷

The correlations between fetal growth and postnatal growth do not imply that the total variability depends mainly from intrauterine growth. Nevertheless, in view of the fact that the observed differences between cohorts defined by birth characteristics are established at an early age (usually in the first months of life), the influence of antenatal events and factors immediately associated with it acquire great importance for understanding the child's growth.

The relationship between birth weight and physical growth is of interest, because neurologic sequelae, suboptimal intelligence and low survival are problems frequently associated to deficient fetal growth and development.³⁸⁻⁴²

VI. OTHER SEQUELAE OF LOW BIRTH WEIGHT

Immediate and late morbidity and sequelae that may result in permanent handicaps are counted among the serious consequences of low birth weight. In pre-term neonates, mortality is fundamentally due to physiologic immaturity, which generates alterations in breathing, in metabolism and in neurological function, as well as in resistance to infections.

Small-for-gestational age infants generally suffer from hypoglycemia and frequently exhibit a similar pathology to that of the pre-term neonate. A chronic deficiency in the quantity and function of the T immunocyte has been described in these infants³⁹ which could explain their greater susceptibility to infection during infancy and preschool age, as the Cauque Study has demonstrated.^{3-9, 27}

It has been described that low birth weight induces sequelae demonstrable at short and long term intervals, which constitute an important cause of defects and disability.⁴⁰⁻⁴² Among these are congenital malformations, early infantile malnutrition, infections, sudden death in infancy, alterations of the central nervous system, mental retardation and behavioral disorders.⁴²

VII. CHANGES IN FETAL GROWTH AND IN NEONATAL MORTALITY

There is great expectation in learning results obtained from the holistic approach in public health to improve problems such as low birth weight. This approach appears to give excellent results judging by changes recently registered in Cuba and Costa Rica.

In 1920, infant mortality in Costa Rica was around 250 per 1,000 livebirths (Table 7). At that time measures to improve the quality of life were beginning to

TABLE 7

INFANT MORTALITY PER 1,000 LIVEBIRTHS IN COSTA RICA

Year	<29 days	29 days - 11 months	Infants
120			250.0
1965	27.2	48.9	76.0
1967	24.3	38.0	62.3
1969	25.4	41.7	67.1
1970	26.2	36.3	61.5
1971	28.7	27.8	56.4
1972	22.8	31.7	54.4
1973	20.8	24.0	44.8
1974	17.7	19.8	37.6
1975	17.7	19.3	37.1

receive particular attention and the result was a progressive increase in the economic income, education and quality of the environment. In the 1950 and 1960 decades infant mortality became stable around figures of 80 per 1,000. The increment in magnitude and coverage of environmental sanitation, nutrition, and medical care programs, as of 1963 ran parallel with a slow descent in infant mortality which then suffered an even more markedly decrease after 1970, coinciding with the strengthening of maternal and child health, sanitation and nutrition programs.

At that time the Rural Health Program (RHP) was implemented taking advantage of the infrastructure created for the control and prevention of malaria in the past decades. The development of the RHP was escalated since 1972, and its goal for 1975 was to cover 600,000 persons of the rural population dispersed in localities of less than 500 inhabitants. This segment represents approximately 30% of the nation's population. At present (1975) a 70% of the dispersed rural population is being covered by nurse auxiliaries and health assistants who are displaced from the Health and Post Centers by jeep, motorcycle, horse, boat or foot.⁴³ Their functions are to carry out census; to vaccinate (tuberculosis, measles, poliomyelitis, smallpox, diphtheria, whooping cough and tetanus) to treat parasitosis (malaria and intestinal parasites), to promote maternal and child health, to promote family planning and nutrition; to treat and refer sick persons; to collaborate in environmental sanitation programs; to give education in hygiene and nutrition, and to participate in community organization efforts.

Data in Table 7 reflect a marked decline in infant mortality as of 1970 when an increase in the level of life, health and development programs was recorded. The change also reflects in part a decrease in neonatal deaths, notorious after 1971 as a result of improvements achieved in the quality of the matreenvironment. Table 8 summarizes the fetal growth situation in a 10-year period at the Institute Materno-Infantil Carit. Already in 1965 the average birth weight was relatively good and the registered frequency of low birth weight was only 110/0. The situation was maintained rather stable up to 1973 when a fall in the frequency of low birth-weight babies was registered reaching a 9.4% value.

The reasons that seem to explain the decrease in neonatal and postneonataj, mortality and in incidence of low birth-weight babies at that date are: a better attentio~ of pregnant women and substantial improvement in family planning. It is quite probable that nutritional changes of secular type which improved the height of the population in a notorious form in the last years⁵ also affected the incidence of low birth-weight neonates. These factors are reflected in a decrease in fetal and maternal mortality observed in previous years (Table 9) and in a decrease in mean parity, which, as illustrated in Tables 7 and 8; were parallel to the reduction in birth rates.

VII. CONTROL AND PREVENTION OF LOW BIRTH WEIGHT

Prevention of low birth weight cannot be achieved through unilateral actions oriented, for example, at control of maternal morbidity or towards the improvement of calories consumption during gestation. The diversity of etiologic factors that intervene, usually in a simultaneous way, in the causality of low birth weight, is sufficient reason for planners of health actions to apply the integral (holistic) approach to its prevention. Table 10 succinctly summarizes different actions that must be implemented within a holistic approach for the prevention of low birth weight.⁴²

Improvements in health and hygiene during childhood are fundamental, particularly in societies where the intergenerational effect of low maternal stature must be reduced. as this reflects a past of malnutrition.}

The practice of optimal reproductive patterns is crucial to diminish the incidence of low birth weight due to premature conception. Measures of this type appear to have had a notorious effect on fetal growth in countries that achieved a marked decrease in mortality in the course of a few years, such as China.⁴⁴ Antenatal care is another important factor for control and prevention of low birth weight of neonates given the high frequency of abnormalities in "high-risk" pregnant women.^{42, 44, 45}

Finally, control and care of the child since early life is fundamental for diagnosis and treatment of present defects and for control and prevention of their sequelae.⁴²⁻⁴⁵

TABLE 9
 BIRTH RATE AND FETAL AND MATERNAL MORTALITY,
 COSTA RICA, 1959-1975

Year	Births per 1,000 population	Mortality per 1,000 live births	
		Maternal	Fetal
1959	48	1.26	20.9
1969	34	1.33	15.2
1973	29	0.93	12.0
1974	30	0.50	12.1
1975	29	*	*

* Pending calculation.

6.

TABLE 10

PREVENTION OF LOW BIRTH WEIGHT AND ITS SEQUELAE

1. *Promotion of health in childhood and adolescence*

Achievement of good nutrition and hygiene
 Prevention of infectious diseases
 Health education on reproduction
 Promotion of good mental health

2. *Practice of the optimal reproductive pattern*

Promotion of conception at optimal ages
 Increment of interval between pregnancies
 Prevention of undesired pregnancies

3. *Antenatal care*

Treatment and prevention of infectious diseases
 Promotion of good nutrition and hygiene
 Education on drugs and stressing agents
 Identification and control of high-risk factors
 Education on perinatal events

4. *Care of neonate and child*

Control of labor and parturition
 Perinatal and neonatal care
 Diagnosis and control of sequelae
 Child care
 Strengthening of infrastructure and recording systems

IX. COMMENT

It is estimated that around 20 million children are born with low birth weight in the world every year; approximately a million and a half of these correspond to Latin America. Low birth weight is the main cause of perinatal mortality and of morbidity and long-term sequelae. Consequently, the control and prevention of low birth weight must constitute one of the higher priorities for national health and development planning programs, particularly in nations that still have high infant mortality rates.

The effects of low birth weight detectable in the field are a high infant mortality (particularly neonatal) and a deficit in physical growth of infants and preschool children.

Studies in a community of the Guatemalan highlands demonstrated that if birth weight is at least 2,750 grams and the child is breast-fed during the first months of life, neonatal survival approaches 100%. The high neonatal mortality observed in underdeveloped areas is therefore due more to the greater number of low birth-weight neonates, than to environmental deficiencies. In effect, neonatal mortality figures observed in the rural community in question, adjusted for weight at birth, were not different from those of an industrial population of a much higher socioeconomic development. Such observation questions the need for creating special resources for the care of premature and fetally malnourished infants in developing countries, in excess of true demands, while it favors the recommendation of measures of preventive nature.

Experience registered in other countries suggests that the holistic approach is the best possible solution for the low birth-weight problem. This approach comprises actions of different nature leading to improve the nutritional status and health of the child and of the future mother, to improve the maternal environment during gestation ... and to favor those circumstances which permit an optimal care of the pregnant and lactating mother and of the neonate and preschool child, for the purpose of guaranteeing a corrective and preventive action of defects and sequelae associated to low birth weight. Such appears to be the reason to explain a achievements observed in Costa Rica and Cuba, where health programs have received priority attention. This position is contrary to the recommendation of isolated measures or interventions whether these are maternal and child health programs or of food supplementation of pregnant mothers, or of any other kind.

A more in-depth analysis of existing information in developing countries is required in regard to benefits derived from different actions and changes in fetal growth. It is also necessary that holistic health and community development plans be structured, comprising actions and recommendations necessary to promote optimal fetal development.

ACKNOWLEDGEMENT

This work was partially financed with funds from the University of Costa Rica, the National Scientific and Technological Research Council (CONICIT), the Ministry of Health, and of the Caja Costarricense del Seguro Social, Costa Rica.

RESUMEN

IMPLICACIONES DE BAJO PESO AL NACER
PARA LA SALUD PUBLICA

El presente estudio interpreta la significación de las características antropométricas del recién nacido de una aldea indígena típica del altiplano de Guatemala y de la población general de Costa Rica. Se ilustra la correlación que existe entre las deficiencias del crecimiento fetal y la mortalidad infantil y el retardo del crecimiento físico. Se discute la significación del retardo del crecimiento fetal en el desarrollo de anomalías y defectos demostrables a corto y a largo plazo.

La información existente indica una alta incidencia de niños con alteraciones de crecimiento y desarrollo fetal en Latinoamérica. Esta circunstancia parece ser la causa principal de la alta mortalidad infantil y del retardo en el desarrollo de muchas poblaciones del Subcontinente,

Algunos países han logrado mejoras substanciales en las condiciones que promueven un adecuado crecimiento fetal mediante la implementación de programas en forma integral (holística). Se proponen las áreas de acción en salud necesarias para lograr un desarrollo fetal adecuado en nuestros países.

BIBLIOGRAPHY

1. Mala, L. J., R. A. Kronmal, J. J. Urrutia & B. Garcia. Antenatal events and postnatal growth and survival of children. Prospective observation in a rural Guatemalan, village. En: *Proceedings Western Hemisphere Nutrition Congress IV*. P. L. White and N. Selvey (Ed.). Mass., Publishing Sciences Group, Inc., p.107,1975.
2. Mata, L. J., J. J. Urrutia, R. A. Kronmal & C. Joplin. Survival and physical growth in infancy and early childhood. *Am. J. Dis. Child.*, 129:561, 1975.
3. Mala, L. J. *The Children of Santa Maria Cauque. A Prospective Study of Health and Growth*. Cambridge, Mass., The MIT Press, 1978,400 p.
4. Dirección General de Estadística y Censos. *Anuarios 1960-1975*. San Jose, Costa Rica.

5. Mat-, L. J. Be E. Mohs. Cambiol culturales y nutricionales en Costa Rica. *Bol. Med. Hosp. Infant. (Mix.)*, 33:579, 1976.
6. Mata, L. J., J. J. Urrutia Be M. Bihar. Infec:don en la mujer embaraz" y en los productos de la concepcion. *Arch. Latinoamer. Nutr.*; 24: 15, 1974.
7. Lechrig, A., J-P. Habicht, G. Guzman & E. de Le6n. MorbiHdad materna y creel- miento fetal en poblaciones rurales de Guatemala. *Arch. Latinoamer. Nutr.*, 22:243,1972.
8. Jurado-Garda, E., A. Abarca, C. Osorio, R. Campos, A. Saavedra, J. Alvare~ & J. Parra. El crecimiento intrauterino. I. Evaluacion del peso y la longitud corporal fetal en la ciudad de Mexico. An'lisis estadistico de 16,807 nacimientos consecutivos de producto unico vivo. *Bol. Med. Hosp. Infant. (Mex.)*, 27:163, 1970.
9. Mata, L. J., J. J. Urrutia, A. caceres Be M. A. Guzman. The biological environ- ment in a Guatemalan rural community. En: *Proceedings Western Hemisphere Nutrition Congress III*. P. L. White and N. Selvey (Eds.). Mount Kisko. N.Y., Futura Publishing Co., Inc., 1972, p. 257.
10. Lubchenco, L. O., D. T. Searls Be J. V. Brazie. Neonatal mortality rate: rela- tionship to birth weight and gestational age. *J. Pediat.*, 81 :814, 1972.
11. Adams, M. S. Be J. D. Niswander. Birth weight of North American Indians: a correction and amplification. *Human Biol.*, 45:351, 1943.
12. Salber, E. J. The significance of birth weight, as illustrated by a comparative study of South African racial groups. *J. Trop. Ped.*, 1:54, 1955.
13. ~ansen, A..A. J. Birth weight, birth length, prematurity and neonatal mortality .. in New Gumeans. *Trop. Geogr. Med.*, 14:341,1962.
14. Sarram, M. & M. Saadatnejadi. Birth weight in Shira (Iran) in relation to mater- nal socioeconomic status. *Obstet. Gynec.*, 30:367, 1967.
15. Banerjee, P. Birth weight of the Bengali newborn: effect of the economic position of the mother. *Ann. Hum. Genet., Lond.* ~33:99,1969.
16. Legg, J., A. M. Davies, R. Prywes, V. V. Sterk & P. Weiskopt. Patterns of low birth weight in West Jerusalem with special reference to maternal origin. *Brit. J. Preo. Soc. Moo.*, 24:89,1970.
17. Coronel, J. G., G. Bustamante & G. Un'be. Prematuridad en nuestro medio. *Pediatria (Rev. Soc. Col. Ped. Puericuli.)*, 10:275, 1968.

18. Luna-JISpe, H., M. Arango, J. Dsu " H. Botero. El peso Y la talla del nacimiento en un grupo de niños de clase económica baja. Manizales, Colombia. *Arch. Latinoamer. Nutr.*, 19:41, 1969.
19. Cor., L., T. F. Pugh, T. H. Inpls" J. E. Gordon. Premature birth as a problem of human populations. *Am. J. Med. Sci.*; 224:343,1952.
20. Erhardt, C. L., G. B. Homi, F. G. Nelson, B. H. Kroll" L. Weiner. Influence of weight and gestational age on perinatal and neonatal mortality by ethnic group. *Am. J. Pub. Health.* 54:1841,1964.

McKeown. T. " J. R. Gibson. Observations on all births (23,970) in Birmingham,1947. IV. "Premature birth". *Brit. Med. J.*, 2:513,1951.
22. Wegman, M. E. International trends in postperinatal mortality. *Am. J. Dis. Child.*, 121:105,1971.
23. US-DHEW. *A Study of Infant Mortality from Linked Records by Birth Weight, Period of Gestation and other Variables*. Washington, D.C., 1972,90 p. (DHEW Publication No. (HSN) 72-1055).
24. Yerushalmy. J. The classification of newborn infants by birth weight and gestational age. *J. Pediat.*, 71: 164, 1967.
25. Tanner, J. M. " A. M. Thomson. Standards for birth weight at gestation periods from 32 to 42 weeks, allowing for maternal height and weight. *Arch. Dis. Childhood*, 45:566, 1970.
26. Lubchenco, L. O., M. Delivoria-Papadopoulos & D. Searls. Long-term follow-up studies of prematurely born infants. II. Influence of birth weight and gestational age on sequelae. *J. Pediat.*, 80:509,1972.

Mata, L. J. Malnutrition-infection interactions in the tropics. *Am. J. Trop. Med. Hyg.*, 24:564,1975.
28. Plank, S. J. & M. C. Milanesi. Infant feeding and infant mortality in rural Chile. *Bull. World Health Org.*, 48:203, 1973.
29. Puffer. R. R. & C. V. Serrano. *Patterns of Mortality in Childhood*. Washington, D. C., Pan American Health Organization, 1973, 470 p. (PAHO Scientific Publication No. 262).
30. Mendez, J. & C. Behrhorst. The anthropometric characteristics of Indian and urban Guatemalans. *HUITInn Bioi.*, 35:457.1963.
31. Habicht, J.P., R. Martorell, C. Yarbrough, R. M. Malina & R. E. Klein. Height and weight standards for preschool children. How relevant are differences in growth potential? *Lancet*, 1:611,1974.

12. Miller, F. J. W., W. A. Billewicz" A. M. Thomson. Growth from birth to adult life on 442 Newcastle-Upon-Tyne children. *Brit. J. Pre». Soc. Med.*, 26:224 1972.
33. Davie, R., N. Butler & H. Goldstein. *From Birth to Seven. The Second Report of the National Child Development Study (1958 Cohort)*. Longman, London, The National Children's Bureau, William Cloves & Sons, 1972, 586 p.
34. Babson, S. G. Growth of low birth-weight infants. *J. Pediat.*, 77:11, 1970.
- H. Fitzhardinge, P. M. & E. M. Steven. The small-for-date infant. I. Later growth-patterns. *Pediatrics*, 49:671, 1972.
36. Morley, D., J. Bicknell" M. Woodland. Factors influencing the growth and nutritional status of infants and young children in a Nigerian village. *Trans. Roy. Soc. Trop. Med. Hyg.*, 62:164,1968.
37. McGregor, I. A., A. K. Rahman, B. Thompson, W. Z. Billewicz" A. M. Thomson. The growth of young children in a Gambian village. *Trans. Roy. Soc. Trop. Med. Hyg.*, 62:341, 1968.
38. Hardy, J. B. Birth weight and subsequent physical and intellectual development. *New Eng. J. Med.*, 289:973, 1973.
39. Chandra, R. K. Immunocompetence in undernutrition. *J. Pediat.*, 81:1194, 1972.
40. Lubchenco, L. O., F. A. Homer, L. H. Reed, I. E. Hix, D. Metcalf, R. Cohig, H. C. Elliott & M. Bourg. Sequelae of premature birth. *Am. J. Dis. Child.*, 106:101,1963.
41. Wright, F. H., R. R. Blough, A. Chamberlin, T. Ernest, W. C. Halstead, P. Meier, R. Y. Moore, R. F. Naunton & F. W. Newell. A controlled follow-up study of small premature born from 1952 through 1956. *Am. J. Dis. Child.*, 124:506, 1972.
42. World Health Organization. *Report on a Meeting on Aetiology, Social Implications and Prevention of Low Birth Weight*. Geneva, 1-5 Sept., 1975. (Unpublished document).
43. Villegas, H. " W. Vargas. *El Progreso de >1100 Rural*. Costa Rica, Ministerio de Salud, 1975.
44. Sidel, V. W. Medical care in the People's Republic of China. *Arch. Int. Med.*, 135:916,1975.

5. Rivcron<ortcguer., R., H. Ferre r-Gard." I. Valdez-Luo. Avances en pediatria y .tendon infantil en Cuba (1959-1974). *Bal. Of. San Pan.*, 80:187, 1976.
46. Chase, H. C. *Relationship of Certain Biologic and Socio-economic Factors to Fetal Infant, and Early Childhood Mortality. 11.Father's Occupation, Infant's Birth Weight and Mother's Age.* Albany, N. V., New York State Dept. of Health, 1962.
47. Jackson, R. L. & G. Kelly. Growth charts for use in pediatric practice.*J. Pediat.*, .27;215,1945,